arXiv:1411.0925v1 [math.PR] 4 Nov 2014

Bounds for a binomial sum involving powers of the summation index

Eliardo G. Costa
Departamento de Estatística, Universidade de São Paulo, São Paulo, Brazil
eliardo@ime.usp.br

November 5, 2014

Abstract

Recently, the properties of a binomial sum related to the multi-link inverted pendulum enumeration problem have been studied. In this note, we establish bounds for this binomial sum.

Keywords: multi-link inverted pendulum, random variable, expectation value.

1 Introduction

The binomial sum $S_p(n)$ defined by

$$S_p(n) = \sum_{j=1}^{n} j^p \binom{n+j}{j}$$

has been studied in the literature because it is related to the multi-link inverted pendulum enumeration problem and it is important to know its properties (see [1] and references therein, for example). In this note, we provide bounds for $S_p(n)$ with p a real positive number using a stochastic approach. Throughout this note $\mathbb{P}(.)$ and $\mathbb{E}[.]$ will denote the probability and expectation operator, respectively.

2 Results

The key of the problem to obtain the bounds is to realize that the sum $S_p(n)$ may be write as a expectation value of a random variable with some adjustments, and using the following Lemma we obtain the result.

Lemma 2.1. *Let $g(x) > 0$ be an even function and nondecreasing on $[0, \infty)$. Suppose that $\mathbb{E}[g(X)] < \infty$. Then for any $x > 0$*

$$\frac{\mathbb{E}[g(X)] - g(x)}{a.s.sup\ g(X)} \leq \mathbb{P}(|X| \geq x) \leq \frac{\mathbb{E}[g(X)]}{g(x)},$$

where $a.s.sup\ g(X) = \inf\{t : \mathbb{P}(g(X) > t) = 0\}$.

Proof. see [5, pg. 52]. □

Theorem 2.1. *For $p > 0$, we have $n^p\binom{2n}{n} \le S_p(n) \le \left[n^p\frac{2n+1}{n+1} + n\right]\binom{2n}{n}$.*

Proof. Consider a random variable X with probability distribution

$$\mathbb{P}(X = j) = c^{-1}\binom{n+j}{j}, \quad j = 0, 1, \ldots, n, \tag{1}$$

where $c = \binom{2n+1}{n}$. It is easy to see that (1) is a probability function since $c = S_0(n) + 1$ (see [2, pg. 159], for example). Then we have

$$\mathbb{P}(|X| \ge n) = \mathbb{P}(X = n) = c^{-1}\binom{2n}{n}.$$

Using Lemma 2.1 with $g(x) = |x|^p$, $p > 0$, we have

$$S_p(n) = c\mathbb{E}[|X|^p] \ge cn^p\mathbb{P}(|X| \ge n) = n^p\binom{2n}{n}.$$

In this case $a.s.sup\ |X|^p = \inf\{t : \mathbb{P}(|X|^p > t) = 0\} = n$. Using Lemma 2.1 we have

$$\mathbb{E}[|X|^p] - n^p \le n\mathbb{P}(|X| \ge n) = nc^{-1}\binom{2n}{n},$$

which implies that $\mathbb{E}[|X|^p] \le n^p + nc^{-1}\binom{2n}{n}$. Then, we obtain

$$S_p(n) = c\mathbb{E}[|X|^p] \le cn^p + n\binom{2n}{n} = n^p\binom{2n+1}{n} + n\binom{2n}{n}.$$

Since $\binom{2n+1}{n} = \frac{2n+1}{n+1}\binom{2n}{n}$ we may write the upper bound as

$$\left[n^p\frac{2n+1}{n+1} + n\right]\binom{2n}{n}.$$

□

Theorem 2.2. *Let $M_p(n) = S_p(n)/[n^p\binom{2n}{n}]$. Then the following statements hold as $n \to \infty$:*

1. *$1 \le M_p(n) \le 2$, if $p > 1$;*
2. *$1 \le M_1(n) \le 3$.*

Proof. Using Theorem 2.1 we have

$$1 \le M_p(n) \le \frac{2n+1}{n+1} + \frac{n}{n^p}.$$

The limit of upper bound is 2 if $p > 1$ and is 3 if $p = 1$, as n approaches infinity.

□

Lemma 2.2. *If $n > 2$, then $\frac{4^n}{2\sqrt{n}} < \binom{2n}{n} < \frac{(2n+2)^n}{(n+1)!}$.*

Proof. see [3, pg. 132] and [4, pg. 297]. □

Corollary 2.1. *If $n > 2$, then $\frac{n^p 4^n}{2\sqrt{n}} < S_p(n) < \left[n^p\frac{2n+1}{n+1} + n\right]\frac{(2n+2)^n}{(n+1)!}$.*

Proof. The result follows from Theorem 2.1 and Lemma 2.2. □

3 Concluding remarks

Theorem 2.1 and Corollary 2.1 provide lower and upper bounds for $S_p(n)$ with p a real positive number. The bounds presented in Corollary 2.1 are less refined than the ones presented in Theorem 2.1, but this result does not present combinatorial numbers. Theorem 2.2 is in agreement with asymptotic results presented in [1].

4 Acknowledgements

The author would like to thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Brazil) and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, Brazil, grant 133211/2011-8) for partial financial support.

References

[1] R. B. Paris and P. J. Larcombe, *On the asymptotic expansion of a binomial sum involving powers of the summation index*, J. Classical Anal. **1** (2), 2012, 113–123.

[2] R. L. Graham, D. E. Knuth, O. Patashnik, *Concrete mathematics*, Addison-Wesley Publishing Company, Reading, 1994, 2nd edition.

[3] W. Sierpiński, *Elementary theory of numbers*, Państwowe Wybawnictwo Naukowe, Warsaw, 1964.

[4] P. Bartoš and Š. Znám, *On symmetric and cyclic means of positive numbers*, Mat.-Fyz. časopis, **16** (3), 1966, 291–298.

[5] Z. Lin and Z. Bai, *Probability inequalities*, Springer and Science Press Beijing, New York, 2010.